

References

1. Mitler MM, Dement WC, Dinges DF. Sleep medicine, public policy, and public health. In: Kryger MH, Roth T, Dement WC, editors. *Principles and practice of sleep medicine*. 3rd ed. Philadelphia: WB Saunders; 2000. p. 580-8.
2. Professional Association of Internes and Residents of Ontario. New call provisions enforceable January 1, 2001. Available: www.pairo.org/about/callschedulingrules.html (accessed 2004 Jan 5).
3. Accreditation Council for Graduate Medical Education. ACGME duty hours standards now in effect for all residency programs. Available: www.acgme.org/Media/news7_1_03.asp (accessed 2004 Jan 5).
4. Gaba DM, Howard SK. Fatigue among clinicians and the safety of patients. *N Engl J Med* 2002;347:1249-55.
5. Rosekind MR, Gregory KB, Miller DL, Co EL, Lebacqz JV. Aircraft accident report: uncontrolled collision with terrain, American International Airways Flight 808, Douglas DC-8, N814CK, U.S. Naval Air Station, Guantanamo Bay, Cuba, August 18, 1993. [Report no. NTSB/AAR-94/04]. Washington, DC: National Transportation Safety Board; 1994.
6. Howard SK, Gaba DM, Rosekind MR, Zarcone VP. The risks and implications of excessive daytime sleepiness in resident physicians. *Acad Med* 2002;77:1019-25.
7. Franzini C. Cardiovascular physiology: the peripheral circulation. In: Kryger MH, Roth T, Dement WC, editors. *Principles and practice of sleep medicine*. 3rd ed. Philadelphia: WB Saunders; 2000. p. 193-203.
8. Parshuram CS, Dhanani S, Kirsh JA, Cox PN. Fellowship training, workload, fatigue and physical stress: a prospective observational study. *CMAJ* 2004;170(6):965-70.

Correspondence to: Dr. David M. Gaba, VA Palo Alto Health Care System, Anesthesia Service, 112A-3801 Miranda Ave., Palo Alto CA 94304; fax 650 852-3423; gaba@stanford.edu

Interventions to promote breast-feeding: applying the evidence in clinical practice

Valerie A. Palda, Jeanne-Marie Guise, C. Nadine Wathen, with the Canadian Task Force on Preventive Health Care

In this document, the Canadian Task Force on Preventive Health Care (CTFPHC) updates its earlier breast-feeding recommendations¹ by presenting evidence on interventions that improve the initiation or duration of breast-feeding (or both). Breast-feeding has been shown in both developing and developed countries to improve the health of infants and their mothers, making it the optimal method of infant nutrition.^{2,3} Although the prevalence of breast-feeding in Canada has risen, with over three-quarters of mothers now initiating breast-feeding, the duration of this practice remains short of the recommended World Health Organization (WHO) targets of exclusive breast-feeding for 6 months and partial breast-feeding for up to 2 years.^{4,5} Recent Canadian data indicate that 22% of recent mothers aged 15-49 years breast-feed for less than 3 months, and 35% do so for at least 3 months.⁶ This premature discontinuation is more a result of difficulty with breast-feeding, including lack of information and support, than of women's choice.⁷ In fact, the number of Canadian hospitals that would qualify as "baby-friendly" according to WHO-UNICEF criteria⁸ was 5 of 523 hospitals responding to a 1993 survey,⁹ and according to UNICEF only a single hospital had that designation in 2002.¹⁰

In a joint endeavour, the CTFPHC and the US Preventive Services Task Force systematically reviewed the randomized trial evidence for the effectiveness of all counselling interventions originating in a clinician's practice (such as antepartum and postpartum support groups, education, telephone support or peer counsellors) to increase the rate of initiation or the duration of breast-feeding.^{11,12} We present here the new CTFPHC recommendations,

based on the joint systematic review as well as a key Canadian trial¹³ published after that review and tailored to the Canadian health care setting (Table 1). Definitions of the levels of evidence and grades of recommendations used in Table 1 are available in an online appendix to this article (www.cmaj.ca/cgi/content/full/170/6/976/DC1).

Interventions consisting of antepartum structured breast-feeding education are effective at improving both initiation and continuation of breast-feeding during the first 2 months postpartum, compared with usual care.¹⁴⁻²⁸ These interventions, consisting of individual or group instruction about breast-feeding knowledge, practical skills and problem-solving techniques, were effective when provided by lactation specialists or nurses, and both single sessions and multiple sessions were effective. Postpartum telephone or in-person support by lactation specialists, nurses or peer counsellors enhanced the effectiveness of these interventions. In addition, the use of peer counsellors improved breast-feeding rates and duration, and these types of programs may represent a cost-effective alternative to professionally delivered services, especially in locations or settings where professional services are scarce or not available.^{13,20,29-32} The CTFPHC recommends against the use of written materials (which have not been shown to be effective when used alone,^{16,19,22,26,33-36} although no harm was demonstrated) and commercial discharge packages (which have been shown to decrease breast-feeding rates).³⁷ Unfortunately, advice from a woman's primary clinician (such as family physician, obstetrician or midwife) has not been sufficiently evaluated, and a research gap remains in this area.

The recommendations presented here (Table 1) do not

Table 1: Canadian Task Force on Preventive Health Care recommendations on interventions to promote breast-feeding

Manoeuvre	Effectiveness	Level of evidence*	Recommendation*
Education programs and postpartum support to promote breast-feeding†	Structured antepartum breast-feeding education improves both initiation and continuation of short-term breast-feeding rates post-partum, compared with usual care‡ In-person or telephone support strengthens the effect of education, leading to an additional 5% to 10% increase in breast-feeding initiation and short-term duration In-person or telephone support by itself may increase both short- and long-term breast-feeding rates	<i>Education</i> Level I — fair ¹⁴⁻¹⁹ Level I — poor ²⁰⁻²⁴ <i>Education + support</i> Level I — fair ^{15,16,19,25-27} Level I — poor ^{20,28}	There is good evidence to recommend provision of structured antepartum educational programs and postpartum support ¹ to promote breast-feeding initiation and duration A recommendation
Peer counselling to promote breast-feeding	Peer counsellors had a significant effect on breast-feeding rates and duration	Level I — fair ¹³ Level I — poor ²⁰ Level II-1 — poor ²⁹⁻³²	There is fair evidence to recommend peer counselling to promote initiation and maintenance of breast-feeding B recommendation
Provision of written materials to new mothers to promote breast-feeding	There is no benefit when written materials are used alone	Level I — good ³³ Level I — fair ^{16,19,26} Level I — poor ^{22,34-36}	There is good evidence to recommend against providing written materials alone to promote breast-feeding D recommendation
Primary health care provider (physician or midwife) advice to expectant or new mothers to promote breast-feeding	Effectiveness is unknown	No studies found	There is insufficient evidence to make a recommendation regarding advice by primary health care providers to promote breast-feeding I recommendation
Provision of commercial discharge packages to new mothers	Women receiving commercial discharge packages had lower breast-feeding rates than patients not receiving packages	Level I (systematic review) — good ³⁷	There is good evidence to recommend against providing commercial discharge packages to new mothers E recommendation
Rooming-in and early maternal contact to promote breast-feeding	The sole new study of rooming-in included multiple interventions, and conclusions could not be drawn New data regarding early maternal contact are insufficient	<i>Rooming in</i> Level I (single study) — fair ³⁸ <i>Early maternal contact</i> Level I (meta-analysis) — good ³⁹⁻⁴²	There is good evidence to recommend rooming-in and early maternal contact to promote breast-feeding A recommendation‡

*Definitions of the levels of evidence and grades of recommendations used by the Canadian Task Force on Preventive Health Care are available in an online appendix to this article (www.cmaj.ca/cgi/content/full/170/6/976/DC1).

†In the studies reviewed, these interventions were usually provided in the clinical setting by lactation specialists or nurses and consisted of individual or group instruction about breast-feeding knowledge, practical skills and problem-solving techniques.

‡The 1994 recommendations of the task force reviewed “good” level I evidence. Those recommendations, which were classified as grade A, are not overturned by the evidence reviewed here.

address the clinical benefits of breast-feeding, which the CTFPHC felt had been established by its earlier breast-feeding recommendations;¹ instead, we have concentrated on recommendations about interventions that change the initiation and duration of breast-feeding.

The Canadian Paediatric Society, Dieticians of Canada and Health Canada have recommended exclusive breast-feeding for at least the first 4 months of life, then continuation of breast-feeding along with complementary foods for up to 2 years and beyond.⁴³ The American Academy of Pediatrics has recommended exclusive breast-feeding for approximately the first 6 months after birth and continued breast-feeding for at least 12 months and thereafter for as long as mutually desired.⁴⁴ Provision of adequate individual and systems-based supports, such as recommended in the

Baby-Friendly Hospital Initiative,⁸ are also recommended by most groups.

This article has been peer reviewed.

From the Department of Medicine, St. Michael's Hospital, Toronto, Ont. (Palda); the Departments of Obstetrics and Gynecology and of Medical Informatics and Outcomes Research, Oregon Health and Science University, Portland, Ore. (Guise); the Offord Centre for Child Studies, McMaster University, Hamilton, Ont. (Wathen)

Competing interests: None declared.

Contributors: Valerie Palda reviewed the evidence and drafted the recommendations and this commentary. Jeanne-Marie Guise and Nadine Wathen reviewed the evidence and draft recommendations, critically revised the current article and reviewed subsequent revisions. The Canadian Task Force on Preventive Health Care critically reviewed the evidence and developed the recommendations according to its methodology and consensus development process. The Canadian Task Force on Preventive Health Care is an independent panel funded by Health Canada.

References

1. Wang EEL. Breastfeeding. In: Canadian Task Force on the Periodic Health Examination. *Canadian guide to clinical preventive health care*. Ottawa: Health Canada; 1994. p. 232-42.
2. Kramer MS, Chalmers B, Hodnett ED, Sevkovskaya Z, Dzvikovich I, Shapiro S, et al. Promotion of Breastfeeding Intervention Trial (PROBIT): a randomized trial in the Republic of Belarus. *JAMA* 2001;285(4):413-20.
3. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet* 2002;360(9328):187-95.
4. World Health Organization. *The optimal duration of exclusive breastfeeding: report of an expert consultation*. Geneva: The Organization; 2002. Available: www.who.int/nut/documents/optimal_duration_of_exc_bfeeding_report_eng.pdf (accessed 2004 Feb 16).
5. World Health Organization. *Global strategy for infant and young child feeding*. Geneva: The Organization; 2003. Available: www.who.int/nut/documents/gis_infant_feeding_text_eng.pdf (accessed 2004 Feb 16).
6. Breastfeeding practices. *Health Indic* [serial online] 2002;2002(1). Ottawa: Statistics Canada; 2002 May. Cat no 82-221-XIE. Available: www.statcan.ca/english/freepub/82-221-XIE/00502/high/canada/cbreast.htm (accessed 2004 Feb 19).
7. Barber CM, Abernathy T, Steinmetz B, Charlebois J. Using a breastfeeding prevalence survey to identify a population for targeted programs. *Can J Public Health* 1997;88(4):242-5.
8. World Health Organization. *Evidence for the ten steps to successful breastfeeding*. Geneva: The Organization; 1998. Available: www.who.int/child-adolescent-health/New_Publications/NUTRITION/WHO_CHD_98.9.pdf (accessed 2004 Feb 16).
9. Dunlop M. Few Canadian hospitals qualify for "Baby Friendly" designation by promoting breast-feeding: survey. *CMAJ* 1995;152(1):87-9.
10. Current status of baby-friendly hospital initiative: March 2002 [table online]. New York: UNICEF; 2002. Available: www.unicef.org/programme/breastfeeding/assets/statusbfhii.pdf (accessed 2004 Feb 16).
11. Guise JM, Palda V, Westhoff C, Chan BKS, Helfand M, Lieu TA. The effectiveness of primary care-based interventions to promote breastfeeding: systematic evidence review and meta-analysis for the US Preventive Services Task Force. *Ann Fam Med* 2003;1:70-8.
12. Palda VA, Guise JM, Wathen CN, Canadian Task Force on Preventive Health Care. Interventions to promote breastfeeding: updated recommendations from the Canadian Task Force on Preventive Health Care. CTFPHC Tech Rep 03-6. London (ON): Canadian Task Force on Preventive Health Care; 2003 Oct.
13. Dennis CL, Hodnett E, Gallor R, Chalmers B. The effect of peer support on breast-feeding duration among primiparous women: a randomized controlled trial. *CMAJ* 2002;166(1):21-8.
14. Duffy EP, Percival P, Kershaw E. Positive effects of an antenatal group teaching session on postnatal nipple pain, nipple trauma and breastfeeding rates. *Midwifery* 1997;13(4):189-96.
15. Pugh LC, Milligan RA. Nursing intervention to increase the duration of breastfeeding. *Appl Nurs Res* 1998;11(4):190-4.
16. Hill PD. Effects of education on breastfeeding success. *Matern Child Nurs J* 1987;16(2):145-56.
17. Kistin N, Benton D, Rao S. Breastfeeding rates among black urban low income women: effect of prenatal education. *Pediatrics* 1990;86:741-6.
18. Brent NB, Redd B, Dworetz A, D'Amico F, Greenberg JJ. Breast-feeding in a low-income population. Program to increase incidence and duration. *Arch Pediatr Adolesc Med* 1995;149(7):798-803.
19. Redman S, Watkins J, Evans L, Lloyd D. Evaluation of an Australian intervention to encourage breastfeeding in primiparous women. *Health Promot Int* 1995;10(2):101-13.
20. Sciacca JP, Dube DA, Phipps BL, Ratliff MI. A breastfeeding education and promotion program: effects on knowledge, attitudes, and support for breastfeeding. *J Community Health* 1995;20(6):473-90.
21. McEnery G, Rao KP. The effectiveness of antenatal education of Pakistani and Indian women living in this country. *Child Care Health Dev* 1986;12(6):385-99.
22. Rossiter JC. The effect of a culture-specific education program to promote breastfeeding among Vietnamese women in Sydney. *Int J Nurs Stud* 1994;31(4):369-79.
23. Wiles LS. The effect of prenatal breastfeeding education on breastfeeding success and maternal perception of the infant. *JOGN Nurs* 1984;13(4):253-7.
24. Reifsnider E, Eckhart D. Prenatal breastfeeding education: its effect on breastfeeding among WIC participants. *J Hum Lact* 1997;13(2):121-5.
25. Oakley A, Rajan L. Social support and pregnancy outcome. *Br J Obstet Gynaecol* 1990;97:155-62.
26. Frank DA, Wirtz SJ, Sorenson JR, Heeren T. Commercial discharge packs and breastfeeding counseling: effects on infant-feeding practices in a randomized trial. *Pediatrics* 1987;80(6):845-54.
27. Serafino-Cross P, Donovan P. Effectiveness of professional breastfeeding home support. *J Nutr Educ* 1992;24(3):117-22.
28. Jones D, West R. Lactation nurse increases duration of breastfeeding. *Arch Dis Child* 1985;60:772-4.
29. Caulfield LE, Gross SM, Bentley ME, Bronner Y, Kessler L, Jensen J, et al. WIC-based interventions to promote breastfeeding among African-American Women in Baltimore: effects on breastfeeding initiation and continuation. *J Hum Lact* 1998;14(1):15-22.
30. Schafer E, Vogel MK, Viegas S, Hausafus C. Volunteer peer counselors increase breastfeeding duration among rural low-income women. *Birth* 1998;25:101-6.
31. Kistin N, Abramson R, Dublin P. Effect of peer counselors on breastfeeding initiation, exclusivity, and duration among low-income urban women. *J Hum Lact* 1994;10(1):11-5.
32. McInnes R, Love J, Stone D. Evaluation of a community-based intervention to increase breastfeeding prevalence. *J Hum Lact* 2000;22:138-45.
33. Curro V, Lanni R, Scipione F, Grimaldi V, Mastroiacovo P. Randomised controlled trial assessing the effectiveness of a booklet on the duration of breastfeeding. *Arch Dis Child* 1997;76(6):500-3.
34. Kaplowitz DD, Olson CM. The effect of an educational program on the decision to breastfeed. *J Nutr Educ* 1983;15:61-5.
35. Loh NR, Kelleher CC, Long S, Loftus BG. Can we increase breast feeding rates? *Ir Med J* 1997;90(3):100-1.
36. Grossman LK, Harter C, Sachs L, Kay A. The effect of postpartum lactation counseling on the duration of breastfeeding in low-income women. *Am J Dis Child* 1990;144(4):471-4.
37. Donnelly A, Snowden H, Renfrew M, Woolridge M. Commercial hospital discharge packs for breastfeeding women [Cochrane review]. In: The Cochrane Library; Issue 3, 2001. Oxford: Update Software.
38. Winikoff B, Myers D, Laukaran VH, Stone R. Overcoming obstacles to breastfeeding in a large municipal hospital: applications of lessons learned. *Pediatrics* 1987;80(3):423-3.
39. De Chateau P, Wiberg B. Long-term effect on mother-infant behaviour of extra contact during the first hour post partum. II. A follow-up at three months. *Acta Paediatr Scand* 1977;66:145-51.
40. Salariya E, Easton P, Cater J. Duration of breastfeeding after early initiation and frequent feeding. *Lancet* 1978;2:1141-3.
41. Thomson M, Hartsock T, Larson C. The importance of immediate postnatal contact: its effect on breastfeeding. *Can Fam Physician* 1979;25:1374-8.
42. Taylor PM, Maloni JA, Taylor FH, Campbell SB. Extra early mother-infant contact and duration of breast-feeding. *Acta Paediatr Scand Suppl* 1985;316:15-22.
43. *Nutrition for healthy term infants. Statement of the Joint Working Group: Canadian Paediatric Society, Dieticians of Canada, Health Canada*. Ottawa: Minister of Health; 1998. Available: www.hc-sc.gc.ca/dca-dea/publications/pdf/infant_e.pdf (accessed 2004 Feb 16).
44. American Academy of Pediatrics, Work Group on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* 1997;100(6):1035-9. Available: aappublications.org/cgi/content/full/pediatrics%3b100/6/1035 (accessed 2004 Feb 16).
45. Harris RP, Helfand M, Woolf SH, Lohr KN, Mulrow CD, Teutsch SM, et al. Current methods of the US Preventive Services Task Force. *Am J Prev Med* 2001;20(3 Suppl):21-35.

Correspondence to: Canadian Task Force on Preventive Health Care, 117-700 Collip Circle, London ON N6G 4X8; fax 519 858-5112; ctf@ctfphc.org

Members of the Canadian Task Force on Preventive Health Care Chair: Dr. John W. Feightner, Professor, Department of Family Medicine, The University of Western Ontario, London, Ont. **Vice-chair:** Dr. Harriet MacMillan, Associate Professor, Departments of Psychiatry and Behavioural Neurosciences and of Pediatrics, Canadian Centre for Studies of Children at Risk, McMaster University, Hamilton, Ont. **Members:** Drs. Paul Bessette, Professeur titulaire, Département d'obstétrique-gynécologie, Université de Sherbrooke, Sherbrooke, Que.; R. Wayne Elford, Professor Emeritus, Department of Family Medicine, University of Calgary, Calgary, Alta.; Denise Feig, Assistant Professor, Department of Medicine, University of Toronto, Toronto, Ont.; Joanne M. Langley, Associate Professor, Departments of Pediatrics and of Community Health and Epidemiology, Dalhousie University, Halifax, NS; Valerie Palda, Assistant Professor, Department of Medicine, University of Toronto, Toronto, Ont.; Christopher Patterson, Professor and Head, Division of Geriatric Medicine, Department of Medicine, McMaster University, Hamilton, Ont.; and Bruce A. Reeder, Professor, Department of Community Health and Epidemiology, University of Saskatchewan, Saskatoon, Sask. **Resource people:** Nadine Wathen, Coordinator; Ruth Walton, Research Associate; and Jana Fear, Research Assistant, Canadian Task Force on Preventive Health Care, Department of Family Medicine, The University of Western Ontario, London, Ont.